



Comparing Spectral Densities of Several Almost Cyclostationary Processes

Mohammad Reza Mahmoudi, Mohsen Maleki, Kirill Borodin,
Kim-Hung Pho and Dumitru Baleanu

EasyChair preprints are intended for rapid dissemination of research results and are integrated with the rest of EasyChair.

April 12, 2022

Comparing Spectral Densities of Several Almost Cyclostationary Processes

Mohammad Reza Mahmoudi^{1,2}, Mohsen Maleki³, Kirill Borodin⁴, Kim-Hung Pho⁵, and Dumitru Baleanu^{6,7}

¹ Institute of Research and Development, Duy Tan University, Da Nang 550000, Vietnam

² Department of Statistics, Faculty of Science, Fasa University, Fasa, Fars, Iran

³ Department of Statistics, Faculty of Science, Shiraz University, Shiraz, Fars, Iran

⁴ Department of Theoretical Physics and Applied Mathematics, Ural Federal University named after the first President of Russia B. N. Yeltsin, Yekaterinburg, Russian Federation

⁵ Fractional Calculus, Optimization and Algebra Research Group, Faculty of Mathematics and Statistics, Ton Duc Thang University, Ho Chi Minh City, Vietnam

⁶ Department of Mathematics, Faculty of Art and Sciences, Cankaya University Balgat 06530, Ankara, Turkey

⁷ Institute of Space Sciences, Magurele-Bucharest, Romania

Abstract. In time series analysis, comparing spectral densities of several processes with almost periodic spectra is an interested problem. The aim of this paper is to give an approach to test the equality among spectral densities of several independent almost periodically correlated (cyclostationary) processes. This approach is based on the limiting distribution for the periodogram and the discrete Fourier transform. The simulation results indicate that the approach well acts.

Key words: Almost periodically correlated processes, Almost cyclostationary processes, Spectral density.

1. Introduction

Comparing spectral densities of several processes is an important topic that has many applications in economics, finance, physics, signal processing, and many others. The researchers like to explore if several time series have the same stochastic mechanism.

Many references have considered the comparison, classification and clustering of two or several time series. For example, De Souza and Thomson (1982), Coates and Diggle (1986), Potscher and Reschenhofer (1988), Diggle and Fisher (1991), Dargahi- Noubary (1992), Diggle and al Wasel (1997), Kakizawa et al. (1998), Timmer et al. (1999), Maharaj [(1999); (2000); (2002); (2005)], Caiado et al. (2006), Eichler (2008), Fokianos and Savvides (2008), Caiado et al. (2009), Dette and Paparoditis (2009), Dette et al. (2010), Dette and Hildebrandt (2011), Jentsch (2012), Jentsch and Pauly (2012), Salcedo et al. (2012), Jentsch and Pauly (2014), Triacca (2016), Mahmoudi et al. (2017) studied these subjects for stationary time series.

But the stationarity assumption is not satisfied in many situations, specially when the processes have a periodic rhythm. In these cases, cyclostationary (CS) and almost cyclostationary (ACS) processes are naturally applied to model the rhythmic component. Mahmoudi et al. (2018b) and Mahmoudi et al. (2018c) considered the comparison, classification and clustering of two CS time series, respectively.

The ACS is a large non-stationary time series class that contained stationary and CS processes. The mean and auto-covariance functions of ACS are almost periodic. The spectra of these processes are supported on lines that are parallel to the main diagonal, $T_j(x) = x \pm \alpha_j, j = 1, 2, \dots$, in spectral square $[0, 2\pi) \times [0, 2\pi)$. The theories and applications of ACS processes have been considered in many resarches such as Gladyshev [(1961); (1963)], Gardner (1991), Hurd (1991), Hurd and Leskow (1992), Leskow and Weron (1992), Gardner (1994), Leskow (1994), Lii and Rosenblatt [(2002); (2006)], Gardner et al. (2006), Hurd and Miamee (2007), Lenart [(2008); (2011)], Napolitano (2012), Lenart (2013), Lenart and Pipien [(2013a), (2013b)], Mahmoudi et al. (2015), Napolitano [(2016a), (2016b)], Mahmoudi and Maleki (2017), Nematollahi et al. (2017), Lenart and Pipien (2017), and Mahmoudi et al. (2018a).

In this work, the asymptotic distribution for the periodogram and discrete Fourier transform of ACS processes will be applied to construct an approach to compare and classify several ACS processes. Section 2 is devoted to notations and preliminaries. The technique to compare and classify the ACS processes is presented in Section 3. The ability of the introduced approach is studied by means of extensive Monte Carlo simulations, and real world problem, in Sections 4 and 5, respectively.

2. Notations and Preliminaries

Definition 1: Almost Periodic Function [Corduneanu (1989)]

A function $f(t): Z \rightarrow R$ is almost periodic if for any $\varepsilon > 0$, there exists a positive integer L_ε such that among any $L_\varepsilon > 0$ consecutive integers there is a positive integer p_ε such that

$$\sup_{t \in Z} |f(t + p_\varepsilon) - f(t)| < \varepsilon.$$

Definition 2: ACS Process [Mahmoudi et al. (2018a)]

A second order process $\{X_t: t \in Z\}$ is called ACS if the process has almost periodic mean, $\mu(t) = E(X_t)$, and autocovariance, $B(t, \tau) = cov(X_t, X_{t+\tau})$, at t , for every $\tau \in Z$.

As Mahmoudi et al. (2018a), the following assumptions have been considered in this work:

- (A1) $\{X_t: t \in Z\}$ is a zero-mean and real-valued time series.
- (A2) X_t is an ACS process.

By these assumptions, the autocovariance function $B(t, \tau)$ can be represented by

$$B(t, \tau) \sim \sum_{\omega \in W_t} a(\omega, \tau) e^{i\omega t},$$

where

$$a(\omega, \tau) = \lim_{n \rightarrow \infty} \left(\frac{1}{n} \sum_{j=1}^n B(j, \tau) e^{-i\omega j} \right),$$

and for fixed τ . Also as Corduneanu (1989) and Hurd (1991) indicated the set $W_\tau = \{\omega \in [0, 2\pi): a(\omega, \tau) \neq 0\}$ is a countable set of frequencies.

(A3) $W = \bigcup_{\tau \in \mathbb{Z}} W_\tau$, is a finite set and the spectra of X_t is supported on lines that are parallel to the main diagonal, $T_j(x) = x \pm \alpha_j, j = 1, 2, \dots$, in spectral square $[0, 2\pi) \times [0, 2\pi)$. Thus we have

$$B(t, \tau) = \sum_{\omega \in W} a(\omega, \tau) e^{i\omega t},$$

and the spectral measure of X_t , will be supported on the set

$$S = \bigcup_{\omega \in W} \{(\nu, \gamma) \in [0, 2\pi)^2: \gamma = \nu - \omega\}.$$

Moreover, the coefficients

$$a(\omega, \tau) = \int_0^{2\pi} e^{i\xi\tau} r_\omega(d\xi),$$

are the Fourier transforms of the measures $r_\omega(\cdot)$.

We note that the r_ω will be identified if the spectral measure of X_t be restricted on the line $\gamma = \nu - \omega$, modulo 2π , where $\omega \in W$.

Remark: In the rest of paper, all equalities of frequencies are modulo 2π .

(A4) r_0 is an absolute continuous measure with respect to the Lebesgue measure.

Dehay and Hurd (1994) shown by considering this assumption and $\sum_{\tau=-\infty}^{\infty} |a(\omega, \tau)| < \infty$, for any $\omega \in W$, result in a spectral density function $f_\omega(\cdot)$ exists such that

$$f_\omega(\nu) = \frac{1}{2\pi} \sum_{\tau=-\infty}^{\infty} a(\omega, \tau) e^{-i\nu\tau}.$$

Consequently, an ACS process with support on a finite number of cyclic frequencies is represented by

$$X_t = \int_0^{2\pi} e^{-itx} \zeta(dx), \quad t \in \mathbb{Z},$$

where ζ is a random spectral measure on $[0, 2\pi)$ such that

$$E(\zeta(d\theta)\overline{\zeta(d\theta')}) = 0, (\theta, \theta') \notin S.$$

As Mahmoudi et al. (2018) indicated, the spectral distribution and density matrices of ζ , are defined by

$$\mathbf{F}(d\lambda) = [F_{k,j}(d\lambda)]_{j,k=1,\dots,m'}$$

and

$$\mathbf{f}(\lambda) = \frac{d\mathbf{F}}{d\lambda} = [f_{k,j}(\lambda)]_{j,k=1,\dots,m'}$$

respectively, where

$$F_{k,j}(d\lambda) = E\left(\zeta(d\lambda + \alpha_k)\overline{\zeta(d\lambda + \alpha_j)}\right), k, j = 1, \dots, m,$$

and $f_{k,j}$ is spectral density correspond to $F_{k,j}$.

Definition 3: Discrete Fourier Transform (DFT)

Assume a sample X_0, \dots, X_{N-1} , from ACS process $\{X_t: t \in Z\}$. The DFT of the finite sequence X_0, \dots, X_{N-1} , is defined by

$$d_X(\lambda) = N^{-1/2} \sum_{t=0}^{N-1} X_t e^{-it\lambda}, \lambda \in [0, 2\pi).$$

Definition 4: Periodogram

Assume that we have a sample X_0, \dots, X_{N-1} , from ACS process $\{X_t: t \in Z\}$. The DFT of the finite sequence X_0, \dots, X_{N-1} , is defined by

$$I_X(\lambda) = |d_X(\lambda)|^2, \lambda \in [0, 2\pi).$$

The distribution of DFT and periodogram of ACS processes are widely studied by Lenart (2013), Lenart and Pipien [(2013a); (2017)] and Mahmoudi et al. (2018a).

3. Methodology

Suppose $\{X_t^{(1)}, t = 1, \dots, n_1\}$, $\{X_t^{(2)}, t = 1, \dots, n_2\}$, ..., $\{X_t^{(l)}, t = 1, \dots, n_l\}$, are l independent ACS processes with m spectral cycles.

Commonly, the researchers want to test the null hypothesis $H_0: X_t^{(1)} \equiv X_t^{(2)} \equiv \dots \equiv X_t^{(l)}$, that is equivalent to $H_0: \mathbf{f}_1 = \mathbf{f}_2 = \dots = \mathbf{f}_l$, such that \mathbf{f}_1, \dots , and \mathbf{f}_l are the spectral density matrices respectively corresponding to $X_t^{(1)}$, ..., and $X_t^{(l)}$. If the null hypothesis H_0 is not accepted then it

can be concluded that at least two time series of the l time series have different rhythms, and if H_0 is accepted consequently the stochastic behaviours of all processes are similar.

Mahmoudi et al. (2018a) introduced the periodogram for ACS time series as

$$\mathbf{I}_X^m(\lambda) = \mathbf{d}_X^m(\lambda) \mathbf{d}_X^{m*}(\lambda),$$

such that

$$\mathbf{d}_X^m(\lambda) = \left(d_X(T_1(\lambda)), d_X(T_2(\lambda)) \dots, d_X(T_m(\lambda)) \right)', \lambda \in [0, 2\pi),$$

where $\mathbf{d}_X^{m*}(\lambda)$ is the complex conjugate transpose of $\mathbf{d}_X^m(\lambda)$.

Lemma 3.1: Let $\{X_t, t \in \mathbb{Z}\}$ is an ACS time series with corresponding spectral density $\mathbf{f}(\lambda)$, $\lambda \in [0, 2\pi)$. Assume $\lambda_1 < \dots < \lambda_J$ are frequencies in $0, 2\pi)$. Then

- (i) $\mathbf{f}(\lambda)$ can be asymptotically estimated by $\hat{\mathbf{f}}(\lambda) = \frac{\mathbf{I}_X^m(\lambda)}{2\pi}$.
- (ii) $\mathbf{d}_X^m(\lambda_j), j = 1, \dots, J$, have the asymptotic and independent m -variate complex normal distributions, $N_m^c\left(0, 2\pi\mathbf{f}(\lambda_j)\right)$.
- (iii) $\mathbf{I}_X^m(\lambda_j), j = 1, \dots, J$, have the asymptotic and independent complex Wishart distributions, $W_m^c\left(\mathbf{1}, 2\pi\mathbf{f}(\lambda_j)\right)$.

Proof: Mahmoudi et al. (2018a). ■

Let $Y_j^{(k)} = \text{Re}\left(\mathbf{d}_{X^{(k)}}^m(\lambda_j)\right), j = 1, \dots, J, k = 1, 2, \dots, l$, and $Z_j^{(k)} = \text{Im}\left(\mathbf{d}_{X^{(k)}}^m(\lambda_j)\right), j = 1, \dots, J, k = 1, 2, \dots, l$, such that $\mathbf{d}_{X^{(k)}}^m(\lambda_j)$ is the $\mathbf{d}_X^m(\lambda_j)$ corresponding to k^{th} time series. As a result of Lemma 3.1, it can be concluded that for $k = 1, 2, \dots, l$, the asymptotic distribution of

$$W_j^{(k)} = \left(Y_j^{(k)}, Z_j^{(k)} \right)' \text{ is } N_{2m}\left(0, \boldsymbol{\Sigma}_j^{(k)}\right), \text{ such that } \boldsymbol{\Sigma}_j^{(k)} = \begin{bmatrix} \mathbf{V}_{Y_j Y_j}^{(k)} & \mathbf{V}_{Y_j Z_j}^{(k)} \\ \mathbf{V}_{Z_j Y_j}^{(k)} & \mathbf{V}_{Z_j Z_j}^{(k)} \end{bmatrix}, \mathbf{V}_{AB} = \text{COV}(A, B).$$

Consequently, the asymptotic distribution of $U^{(k)} = \sum_{j=1}^J W_j^{(k)}$ is $N_{2m}\left(0, \boldsymbol{\Sigma}^{(k)}\right)$, such that $\boldsymbol{\Sigma}^{(k)} = \boldsymbol{\Sigma}_1^{(k)} + \dots + \boldsymbol{\Sigma}_J^{(k)}$.

3.1. Test of Hypothesis

As previous discussion, usually, the researchers want to test the null hypothesis $H_0: \mathbf{f}_1 = \mathbf{f}_2 = \dots = \mathbf{f}_l$, that is equivalent to $H_0: \boldsymbol{\Sigma}^{(1)} = \boldsymbol{\Sigma}^{(2)} = \dots = \boldsymbol{\Sigma}^{(l)}$. It can be concluded that the

asymptotic distribution of $U = U^{(1)} + U^{(2)} + \dots + U^{(l)}$ is $N_{2m}(0, \Sigma)$, such that $\Sigma = \Sigma^{(1)} + \Sigma^{(2)} + \dots + \Sigma^{(l)}$.

Therefore the asymptotic distribution of statistic

$$\chi^2 = (U)'(\Sigma)^{-1}(U),$$

is chi-square with $2m$ degrees of freedom, $\chi^2(2m)$.

The asymptotic distribution of statistic χ^2 can be applied to establish test of hypothesis about the null hypothesis H_0 . As can be seen, the statistic χ^2 is related to the unknown parameter Σ .

Remark 3.1: To produce a sample of size N_i for the discrete Fourier transform of i^{th} process, different bootstrap approaches can be applied. In this research we used the moving block bootstrap (MBB) (more details are given in Synowiecki (2007)).

Let $\mathbf{S} = \frac{(N_1-1)\mathbf{S}^{(1)} + (N_2-1)\mathbf{S}^{(2)} + \dots + (N_l-1)\mathbf{S}^{(l)}}{N_1 + N_2 + \dots + N_l - l}$, as the pooled covariance matrix of $\mathbf{S}^{(1)}, \dots, \mathbf{S}^{(l)}$, such

that $\mathbf{S}^{(k)} = \mathbf{S}_1^{(k)} + \dots + \mathbf{S}_j^{(k)}$, $\mathbf{S}_j^{(k)} = \begin{bmatrix} \widehat{\mathbf{V}}_{Y_j Y_j}^{(k)} & \widehat{\mathbf{V}}_{Y_j Z_j}^{(k)} \\ \widehat{\mathbf{V}}_{Z_j Y_j}^{(k)} & \widehat{\mathbf{V}}_{Z_j Z_j}^{(k)} \end{bmatrix}$, and $\widehat{\mathbf{V}}_{AB} = \widehat{COV}(A, B)$. If the null

hypothesis $H_0: \Sigma^{(1)} = \Sigma^{(2)} = \dots = \Sigma^{(l)}$, be true, then the covariance matrix Σ can be consistently estimated by \mathbf{S} , and consequently by using the WLLN (Weak Law of Large Numbers), the asymptotic distribution of the test statistic

$$\chi^{2*} = (U)'(\mathbf{S})^{-1}(U),$$

is $\chi^2(2m)$. Therefore, for a given size α , H_0 is rejected if $\chi^{2*} > \chi_{\alpha}^2(2m)$.

4. Simulation Study

To analyze the accuracy of proposed method, we generated

$$(n_1, n_2, n_3) = \{(100, 50, 75), (150, 75, 100), (200, 150, 100), (500, 250, 300)\}$$

observations from the ACS processes

$$X_t^{(i)} = (1 + \cos(\omega_i t))Y_t^{(i)}, \omega \in (0, \infty), i = 1, 2, 3,$$

where

$$Y_t^{(i)} = Z_t^{(i)} + 0.5Z_{t-1}^{(i)},$$

and $Z_t^{(i)}, i = 1, 2, 3$, are independent sequences of IIDN(0,1).

The spectral mass of $X_t^{(i)}$ is supported on the lines given by

$$T_1(x) = x, T_2(x) = x + \omega_i, T_3(x) = x - \omega_i, T_4(x) = x - 2\omega_i, T_5(x) = x + 2\omega_i.$$

Figure 1 indicates the spectral plane $[0, 2\pi)^2$, for

$$\omega_i = \{0.75, 1.5, 2.25\}.$$




Figure 1: The spectral square of the process, Left: $\omega_i = 0.5$, Middle: $\omega_i = 1$, and Right: $\omega_i = 2$.

First, we estimated the Type I error probability ($\hat{\alpha}$) and power ($\hat{\pi}$) based on 1000 replications and 1000 iterations. Then we graph Q-Q plot for the test statistic χ^2^* based on the computed values of the simulation runs.

Table 1 report the values of $\hat{\alpha}$ (in rows: 1th, 14th and 27th) and $\hat{\pi}$ (other rows). The results indicates that the values of $\hat{\alpha}$ is very close to the considered size ($\alpha = 0.05$), especially when (n_1, n_2, n_3) grows. Also the power studies show that the proposed method excellently discriminate H_0 from H_1 .

Table 1: The values of $\hat{\alpha}$ and $\hat{\pi}$ for the introduced approach

			(n_1, n_2, n_3)			
ω_1	ω_2	ω_3	(100, 50, 75)	(150, 75, 100)	(200, 150, 100)	(500, 250, 300)
0.75	0.75	0.75	0.052	0.050	0.050	0.049
		1.5	0.736	0.845	0.909	0.976
		2.25	0.727	0.828	0.925	0.978
	1.5	0.75	0.706	0.837	0.907	0.999
		1.5	0.760	0.864	0.959	0.994
		2.25	0.766	0.866	0.955	0.987
	2.25	0.75	0.718	0.812	0.903	0.970
		1.5	0.758	0.889	0.969	0.985
		2.25	0.800	0.883	0.945	0.986
1.5	0.75	0.75	0.772	0.864	0.969	1.000
		1.5	0.727	0.842	0.939	0.972
		2.25	0.760	0.861	0.946	0.989
	1.5	0.75	0.741	0.842	0.917	0.996
		1.5	0.051	0.051	0.050	0.050
		2.25	0.725	0.845	0.906	0.970

	2.25	0.75	0.782	0.897	0.963	0.983
		1.5	0.717	0.835	0.919	0.993
		2.25	0.790	0.880	0.943	0.997
2.25	0.75	0.75	0.791	0.888	0.952	0.984
		1.5	0.777	0.857	0.963	0.999
		2	0.729	0.847	0.904	0.994
	1.5	0.75	0.779	0.870	0.965	0.989
		1.5	0.797	0.871	0.959	0.989
		2.25	0.718	0.835	0.933	0.978
	2.25	0.75	0.718	0.844	0.904	0.976
		1.5	0.724	0.844	0.933	0.973
		2.25	0.052	0.051	0.049	0.049

5. Real Data

This section is devoted to illustrate the ability of introduced approach in practical cases. The dataset includes the first difference of centered moving average filter 2×12 moving average (MA) applied for logarithm of industrial production index (IPI) in Poland (2005 = 100%) since January 1995 until December 2009, Lenart and Pipien (2013b). We split this dataset in three parts with equal sizes. The spectral frequency squares of these parts are given in Figure 2. The results detect ACS time series with spectra on the lines $T_j(x) = x \pm \alpha$, $\alpha \in \{0.062, 0.153, 0.258\}$. This result verifies the given result in Lenart and Pipien (2013b). Then the introduced technique is used to test the hypothesis $\Sigma^{(1)} = \Sigma^{(2)} = \Sigma^{(3)}$ (or equivalently, $\mathbf{f}_1 = \mathbf{f}_2 = \mathbf{f}_3$). Table 2 summarizes the results. As can be seen, since the p value is more than 0.05, thus the null hypothesis can not be rejected and consequently the stochastic behaviours of all processes are similar.




Figure 2: Spectral frequency square (Left: Part 1, Middle: Part 2, Right: Part 3)

Table 2: Testing the equality of different parts

Test Statistic	P-Value
----------------	---------

$\chi^2^* = 4.193$	0.651
--------------------	-------

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- Caiado, J., Crato, N., Pena, D. (2006). A Periodogram-based Metric for Time Series Classification, *Computational Statistics & Data Analysis*, **50**, 2668-2684.
- Caiado, J., Crato, N., Pena, D. (2009). Comparison of times series with unequal length in the frequency domain, *Comm. Statist. Simulation Comput.*, **38**, 527–540.
- Coates, D. S., Diggle, P. J. (1986). Tests for Comparing Two Estimated Spectral Densities, *Journal of Time Series Analysis*, **7**, 7-20.
- Corduneanu, C. (1989). *Almost Periodic Functions*, Chelsea: New York.
- Dargahi-Noubary, G. R. (1992). Discrimination between Gaussian Time Series based on Their Spectral Differences, *Communications in Statistics (Theory and Methods)*, **21**, 2439-2458.
- Dehay, D., Hurd, H. (1994). Representation and estimation for periodically and almost periodically correlated random processes. In: W.A. Gardner (Ed.), *Cyclostationarity in Communications and Signal Processing*, IEEE Press, 295–329.
- De Souza, P., Thomson, P. (1982). Lpc distance measures and statistical tests with particular reference to likelihood ratio, *IEEE Transactions Accoustics Speech Signal Process*, **30**, 304-315.
- Dette, H., Hildebrandt, T. (2011). A note on testing hypothesis for stationary processes in the frequency domain, *J. Multivariate Anal.*, **104**, 101–114.
- Dette, H., Kinsvater, T., and Vetter, M. (2010). Testing nonparametric hypotheses for stationary processes by estimating minimal distances, *Journal of Time Series Analysis*, **32**, 447-461.
- Dette, H., Paparoditis, E. (2009). Bootstrapping frequency domain tests in multivariate time series with an application to comparing spectral densities, *Journal of the Royal Statistical Society B*, **71**, 831-8571.
- Diggle, P. J., Fisher, N. I. (1991). Nonparametric Comparison of Cumulative Periodograms, *Applied Statistics*, **40**, 423-434.
- Diggle, P. J., al Wasel, I. (1997). Spectral Analysis of Replicated Biomedical Time Series, *Applied Statistics*, **46**, 31-71.
- Eichler, M. (2008). Testing nonparametric and semiparametric hypotheses in vector stationary processes, *J. Multivariate Anal.*, **99** 968–1009.
- Fokianos, K., Savvides, A. (2008). On Comparing Several Spectral Densities. *Technometrics*, **50 (3)**, 317-331.

- Gardner, W. A. (1991). Exploitation of Spectral Redundancy in Cyclostationary Signals. *IEEE Signal Processing Magazine*, **8** (2), 14–36.
- Gardner, W. A., ed. (1994). *Cyclostationarity in Communications and Signal Processing*. IEEE Press, New York.
- Gardner, W. A., Napolitano, A., Paura, L. (2006). Cyclostationarity: Half a Century of Research, *Signal Processing*, **86**, 639-697.
- Gladyshev, E. G. (1961). Periodically Correlated Random Sequences. *Soviet Math. Dokl.*, **2**, 385-388.
- Gladyshev, E. G. (1963). Periodically and Almost Periodically Correlated Random Processes with a Continuous Time Parameter. *Theory Probab. Appl.*, **8**, 173–177.
- Hurd, H. (1991). Correlation theory of almost periodically correlated processes. *J. Multivariate Anal.*, **37**, 24–45.
- Hurd, H., Leskow, J. (1992). Strongly Consistent and Asymptotically Normal Estimation of the Covariance for Almost Periodically Correlated Processes. *Statist. Decisions*, **10**, 201–225.
- Hurd, H. L., Miamee, A. G. (2007). *Periodically Correlated Random Sequences: Spectral Theory and Practice*. Wiley, Hoboken.
- Jentsch, C. (2012). A new frequency domain approach of testing for covariance stationarity and for periodic stationarity in multivariate linear processes, *J. Time Series Anal.*, **33**, 177–192.
- Jentsch, C., Pauly, M. (2012). A note on using periodogram-based distances for comparing spectral densities, *Statist. Probab. Lett.*, **82**, 158–164.
- Jentsch, C., Pauly, M. (2015). Testing equality of spectral densities using randomization techniques, *Bernoulli*, **21**(2), 697–739.
- Kakizawa, Y., Shumway, R. H., Taniguchi, M. (1998). Discrimination and Clustering for Multivariate Time series, *Journal of the American Statistical Association* **93**, 328-340.
- Lenart, L. (2008). Asymptotic properties of periodogram for almost periodically correlated time series. *Probability and Mathematical Statistics*. **28** (2), 305-324.
- Lenart, L. (2011). Asymptotic distributions and subsampling in spectral analysis for almost periodically correlated time series. *Bernoulli*, **17**(1): 290–319.
- Lenart, L. (2013). Non-parametric Frequency Identification and Estimation in Mean for Almost Periodically Correlated Time Series, *Journal of Multivariate Analysis*, **115**, 252-269.
- Lenart L., Pipien, M. (2013a). Seasonality Revisited - Statistical Testing for Almost Periodically Correlated Processes, *Central European Journal of Economic Modelling and Econometrics*, **5**, 85-102.
- Lenart, L., Pipien, M. (2013b). Almost Periodically Correlated Time Series in Business Fluctuations Analysis, *Acta Physica Polonica A*, **123**(3), 567-583.
- Lenart, L., Pipien, M. (2017). Non-Parametric Test for the Existence of the Common Deterministic Cycle: The Case of the Selected European Countries, *Central European Journal of Economic Modeling and Econometrics*, **9** (3), 201-241.
- Leskow, J. (1994). Asymptotic normality of the spectral density estimator for almost periodically correlated stochastic processes. *Stoch. Process. Appl.*, **52**, 351–360.
- Leskow, J., Weron, A. (1992). Ergodic Behavior and Estimation for Periodically Correlated Processes. *Statist. Probab. Lett.*, **15**, 299–304.

- Lii, K.-S., Rosenblatt, M. (2002). Spectral Analysis for Harmonizable Processes. *Ann. Statist.*, **30** (1), 258–297.
- Lii, K.-S., Rosenblatt, M. (2006). Estimation for Almost Periodic Processes. *Ann. Statist.*, **34** (3), 1115–1139.
- Maharaj, E.A. (1999). Comparison and classification of stationary multivariate time series, *Pattern Recognition*, **32**, 1129–1138.
- Maharaj, E.A. (2000). Clusters of time series, *Journal of Classification*, **17**, 297–314.
- Maharaj, E. A. (2002). Comparison of Non-stationary Time Series in the Frequency Domain, *Computational Statistics & Data Analysis*, **40**, 131-141.
- Maharaj, E.A. (2005). Using wavelets to compare time series patterns, *International Journal of Wavelets, Multiresolution and Information Processing*, **3**, 511–521.
- Mahmoudi, M. R., Heydari, M. H., Avazzadeh, Z. (2018a). On the Asymptotic Distribution for the Periodograms of Almost Periodically Correlated (Cyclostationary) Processes, *Digital Signal Processing*, **81**, 186-197.
- Mahmoudi, M. R., Heydari, M. H., Avazzadeh, Z. (2018b). Testing the Difference between Spectral Densities of Two Independent Periodically Correlated (Cyclostationary) Time Series Models, *Communications in Statistics—Theory and Methods*, In Print.
- Mahmoudi, M. R., Heydari, M. H., Roohi, R. (2018c). A new method to compare the spectral densities of two independent periodically correlated time series, *Mathematics and Computers in Simulation*, DOI: 10.1016/j.matcom.2018.12.008.
- Mahmoudi, M. R., Maleki, M. (2017). A New Method to Detect Periodically Correlated Structure, *Computational Statistics*, **32** (4), 1569-1581.
- Mahmoudi, M. R., Maleki, M., Pak, A. (2017). Testing the Difference between Two Independent Time Series Models. *Iranian Journal of Science and Technology: Sciences* **41**(3): 665-669.
- Mahmoudi, M. R., Nematollahi, A. R., Soltani, A. R. (2015). On the Detection and Estimation of Simple Processes, *Iranian Journal of Science and Technology*, **39** A2, 239-242.
- Napolitano, A. (2012). *Generalizations of Cyclostationary Signal Processing: Spectral Analysis and Applications*, Wiley-IEEE Press.
- Napolitano, A. (2016a). Cyclostationarity: Limits and generalizations, *Signal Processing*, **120**, 323-347.
- Napolitano, A. (2016b). Cyclostationarity: New trends and applications, *Signal Processing*, **120**, 385-408.
- Nematollahi, A. R., Soltani, A. R., Mahmoudi, M. R. (2017). Periodically Correlated Modeling by Means of the Periodograms Asymptotic Distributions, *Statistical Papers*, **58** (4), 1267-1278.
- Potscher, B., Reschenhofer, E. (1988). Discriminating between two spectral densities in case of replicated observations, *Journal of Time Series Analysis*, **9**, 221-224.
- Salcedo, G. E., Porto, R. F., Morettin, P. A. (2012). Comparing non-stationary and irregularly spaced time series, *Computational Statistics & Data Analysis*, **56**(12), 3921–3934.
- Synowiecki, R. (2007). Consistency and Application of Moving Block Bootstrap for Nonstationary Time Series with Periodic and Almost Periodic Structure, *Bernoulli*, **13**, 1151–1178.

Timmer, J., Lauk, M., Vach, W., Lucking, C.H. (1999). A test for the difference between spectral peak frequencies, *Computational Statistics & Data Analysis*, **30**, 45–55.

Triacca, U. (2016). Measuring the Distance between Sets of ARMA Models, *Econometrics*, **4(3)**, 32.

Samadianfard, Saeed, et al. "Wind speed prediction using a hybrid model of the multi-layer perceptron and whale optimization algorithm." *Energy Reports* 6 (2020): 1147-1159.

Taherei Ghazvinei, Pezhman, et al. "Sugarcane growth prediction based on meteorological parameters using extreme learning machine and artificial neural network." *Engineering Applications of Computational Fluid Mechanics* 12.1 (2018): 738-749.

Qasem, Sultan Noman, et al. "Estimating daily dew point temperature using machine learning algorithms." *Water* 11.3 (2019): 582.

Mosavi, Amir, and Atieh Vaezipour. "Reactive search optimization; application to multiobjective optimization problems." *Applied Mathematics* 3.10A (2012): 1572-1582.

Shabani, Sevda, et al. "Modeling pan evaporation using Gaussian process regression K-nearest neighbors random forest and support vector machines; comparative analysis." *Atmosphere* 11.1 (2020): 66.

Ghalandari, Mohammad, et al. "Aeromechanical optimization of first row compressor test stand blades using a hybrid machine learning model of genetic algorithm, artificial neural networks and design of experiments." *Engineering Applications of Computational Fluid Mechanics* 13.1 (2019): 892-904.

Mosavi, Amir. "Multiple criteria decision-making preprocessing using data mining tools." *arXiv preprint arXiv:1004.3258* (2010).

Karballaezadeh, Nader, et al. "Prediction of remaining service life of pavement using an optimized support vector machine (case study of Semnan–Firuzkuh road)." *Engineering Applications of Computational Fluid Mechanics* 13.1 (2019): 188-198.

Asadi, Esmail, et al. "Groundwater quality assessment for sustainable drinking and irrigation." *Sustainability* 12.1 (2019): 177.

Mosavi, Amir, and Abdullah Bahmani. "Energy consumption prediction using machine learning; a review." (2019).

Dineva, Adrienn, et al. "Review of soft computing models in design and control of rotating electrical machines." *Energies* 12.6 (2019): 1049.

Mosavi, Amir, and Timon Rabczuk. "Learning and intelligent optimization for material design innovation." In *International Conference on Learning and Intelligent Optimization*, pp. 358-363. Springer, Cham, 2017.

Torabi, Mehrnoosh, et al. "A hybrid machine learning approach for daily prediction of solar radiation." *International Conference on Global Research and Education*. Springer, Cham, 2018.

Mosavi, Amirhosein, et al. "Comprehensive review of deep reinforcement learning methods and applications in economics." *Mathematics* 8.10 (2020): 1640.

Ahmadi, Mohammad Hossein, et al. "Evaluation of electrical efficiency of photovoltaic thermal solar collector." *Engineering Applications of Computational Fluid Mechanics* 14.1 (2020): 545-565.

Ghalandari, Mohammad, et al. "Flutter speed estimation using presented differential quadrature method formulation." *Engineering Applications of Computational Fluid Mechanics* 13.1 (2019): 804-810.

Ijadi Maghsoodi, Abteen, et al. "Renewable energy technology selection problem using integrated h-swara-multimoor approach." *Sustainability* 10.12 (2018): 4481.

Mohammadzadeh S, Danial, et al. "Prediction of compression index of fine-grained soils using a gene expression programming model." *Infrastructures* 4.2 (2019): 26.

Sadeghzadeh, Milad, et al. "Prediction of thermo-physical properties of TiO₂-Al₂O₃/water nanoparticles by using artificial neural network." *Nanomaterials* 10.4 (2020): 697.

Choubin, Bahram, et al. "Earth fissure hazard prediction using machine learning models." *Environmental research* 179 (2019): 108770.

Emadi, Mostafa, et al. "Predicting and mapping of soil organic carbon using machine learning algorithms in Northern Iran." *Remote Sensing* 12.14 (2020): 2234.

Shamshirband, Shahaboddin, et al. "Developing an ANFIS-PSO model to predict mercury emissions in combustion flue gases." *Mathematics* 7.10 (2019): 965.

Salcedo-Sanz, Sancho, et al. "Machine learning information fusion in Earth observation: A comprehensive review of methods, applications and data sources." *Information Fusion* 63 (2020): 256-272.